

# **Summary of Forum 2008 Findings on Microlensing**

Scott Gaudi (OSU)



Andy Boden (MSC)

J.P. Beaulieu (IAP), Dave Bennett (UND), Edward Cheng  
(Conceptual Analytics), Kem Cook (LLNL), Andy Gould  
(OSU), John Mather (NASA), Charley Noecker (Ball),  
Domenick Tenerelli (Lockheed)

April 21, 2009 Missions for Exoplanets

# Microlensing Peculiarities


- Microlensing planet community is comparatively small and collaborative.
- General consensus (and movement) on forward directions.
- Two primary paths forward
  - Ground based, 1-5 years
    - Frequency of planets  $>M_{\oplus}$  beyond the snow line.
  - Space based, 5-10 years,  $\sim \$300\text{M}$ 
    - Complete census of planets with mass greater than Mars and  $a > 0.5$  AU, including habitable planets and free floating planets.



## Beyond the snow line:


- Location of giant planet formation (and our giant planets).
- ‘Failed Jupiters’
- Source of water

*Ground-based  $\mu$ lensing surveys probe planets with  $M > M_{\oplus}$  beyond the snow-line.*



*A space-based survey will provide a complete census of planetary systems with mass greater than Mars and  $a > 0.5$  AU (from 0 to  $\infty$  with Kepler), including habitable planets.*

# Detecting Planets



Primary event:


- Smooth, symmetric
- Typically 20 days

Planetary perturbation to images

- Short-timescale bump
- Measure:
  - Projected Separation
  - Mass Ratio

$$t_p = q^{1/2} t_E \approx 1 \text{ day} \left( \frac{M_p}{M_J} \right)^{1/2}$$


# Microlensing is directly sensitive to planet mass



- Works by perturbing images
- Does not require light from the lens or planet.
- Sensitive to planets in the disk and bulge with  $D_{OL}=1-8$  kpc
- Most sensitive to planets near the Einstein radius
- Sensitive to wide or free-floating planets
- Not sensitive to very close planets

# Very Low Mass Planets

Signal magnitude is *independent* of planet mass.




- Magnitude depends on separation of planet from image.
- Duration depends on mass.

$$t_p = q^{1/2} t_E \approx 2 \text{ hrs} \left( \frac{M_p}{M_\oplus} \right)^{1/2}$$

- Signals get rarer and briefer.
- Detection Probability  
~ few %
- **Large** signals for low-mass (Earth-mass) planets
- **Sub Mars-mass planets detectable**

# Sensitivity Depends Weakly on Host Mass



Sensitive to planets around:

- Main-sequence stars with  $M < M_\odot$
- Brown dwarfs
- Remnants




# Microlensing Strengths

- Peak sensitivity beyond the snow line.
  - 50-100K
- Sensitivity down to *very* low-mass planets.
  - Mass greater than that of  $\sim 10\%$  Mars.
- Sensitivity to long-period and free-floating planets.
  - 0.5 AU -  $\infty$
- Sensitivity to planets over a wide range of host masses.
  - $M < M_{\odot}$
- Sensitivity to planets throughout the Galaxy.
  - 1-8 kpc
- Sensitivity to multiple-planet systems.

# Commonly Heard Complaints...

- But you don't know anything about the star, orbits, etc!
  - Typically can measure host star and planet masses to  $\sim 10\text{-}20\%$ .
  - In some special cases can learn something about the orbit.
- But the systems are so far away and faint!
  - Sufficiently bright to measure flux, color, and in some cases get spectra.
- But you only see it once!
  - Signals are large and unambiguous.
- Demographics of planetary systems.


OGLE-2005-BLG-071  
(Udalski et al 2005)



$$M_p \sim 3.5 M_J, \quad r \sim 3.6 \text{ AU}$$

$$M_* \sim 0.46 M_\odot, \quad D_{OL} \sim 3.3 \text{ kpc}$$


OGLE-2004-BLG-235  
MOA-2004-BLG-53  
(Bond et al 2004)



$$M_p \approx 2.5 M_J, \quad r \approx 4.3 \text{ AU}$$


$$M_* \sim 0.65 M_\odot, \quad D_{OL} \sim 6.5 \text{ kpc}$$

# First Four Detections.



Two Jovian-mass planets  
Two “Super Earth” planets


OGLE-2005-BLG-390  
(Beaulieu et al 2006)



$$M_p \sim 5.5 M_\oplus, \quad r \sim 2.6 \text{ AU}$$

$$M_* \sim 0.22 M_\odot, \quad D_{OL} \sim 6.6 \text{ kpc}$$

OGLE-2005-BLG-169  
(Gould et al 2006)



$$M_p \sim 13 M_\oplus, \quad r \sim 3.5 \text{ AU}$$

$$M_* = 0.5 M_\odot, \quad D_{OL} = 2.7 \text{ kpc}$$

# Cool “Super Earths” Are Common.

Two low-mass detections imply:

**~20% of stars have Super Earths between 1.6-4.3 AU  
(10-30%)**


**$dN/d\log a \sim 0.5$  at  $\sim 3$  AU (0.4-1.6 at 90%)**

**$dN/d\log a \sim 0.3$  at  $\sim 0.15$  AU (Mayor et al 2009, HARPS)**

Also:

**Cool Super Earths are more common than cool Jupiters**

# Multiple System



- High-magnification Event
  - μFUN, OGLE, MOA
- Must include two planets, finite source, orbital motion, and parallax
- Yields full star and planet masses, information on orbital speed of Saturn and inclination!

A  $\sim 0.5 M_{\odot}$  late K-dwarf at  $\sim 1.5$  kpc

Finite  
Source

$$\theta_E \cong 1.48 \text{ mas}$$

Parallax

$$\tilde{r}_E \cong 2.76 \text{ AU}$$

AO Imaging



$$D_l \cong 1.49 \pm 0.13 \text{ kpc}$$

$$M = 0.50 \pm 0.05 M_{\odot}$$

# The OGLE-2006-BLG-109L Planetary System

## Planet b:

$$\text{Mass} = 0.71 \pm 0.08 M_{\text{Jup}}$$


$$\text{Semimajor Axis} = 2.3 \pm 0.2 \text{ AU}$$

## Planet c:

$$\text{Mass} = 0.27 \pm 0.03 M_{\text{Jup}} = 0.90 M_{\text{Sat}}$$

$$\text{Semimajor Axis} = 4.6 \pm 0.5 \text{ AU}$$


# Analog of Jupiter/Saturn






# Low Mass Planet Orbiting a Substellar Host

(Bennett et al 2008)




$$M = 0.06 \pm 0.03 M_{\odot}$$

$$m = 3.3^{+4.9}_{-1.6} M_{\oplus}$$




# The Most Massive M Dwarf Planet?

(Dong et al 2008)



Dong et al. 2008



$$M = 0.46 \pm 0.04 M_{\odot}$$

$$D_l = 3.3 \pm 0.4 \text{ kpc}$$


$$v_{\text{LSR}} = 103 \pm 14 \text{ km s}^{-1}$$

$$m = 3.5 \pm 0.3 M_{\text{Jup}}$$


$$r_{\perp} = 3.6 \pm 0.2 \text{ AU}$$

$$T_{eq} \sim 50 \text{ K}$$


More planets.




2008: Jupiter-mass planet.



2007: “Buried” Jupiter-mass planet, exhumed by Dong et al. (2008)



2008: “Buried” Saturn-mass planet, likely in the bulge.



2008: Brown dwarf companion?

# More on the Way...

- **Eight likely planets discovered in the 2007-2008 seasons.**
- **14 likely planets discovered to date, in 12 systems.**
- **2009 season underway (possible detection of a Jupiter-mass planet by MOA)**
- **Can expect  $\sim 4$  planets per year.**



# What's Next?

- **Current setup (alert/follow-up) saturated**
  - Nearly all of the useable bulge monitored
  - Many events cannot be monitored
  - Monitoring one event at a time too inefficient
- **A new strategy**
  - Dispense with alert/follow-up
  - Simultaneously detect and monitor microlensing events

# What is Required?

- Event Rate

- Primary Event Rate


$$\Gamma \approx 10^{-5} \text{ yr}^{-1}$$

- Detection Probability

$$P \approx A_0 \theta_p \approx 1\% \left( \frac{M_p}{M_{\text{Earth}}} \right)^{1/2}$$

- Detections Per Year

$$N \approx n_F \Omega \Phi \Gamma P \approx 10 \text{ yr}^{-1} \left( \frac{\Omega}{10 \square^\circ} \right) \left( \frac{\Phi}{10^7 / \square^\circ} \right) \left( \frac{\Gamma}{10^{-5} \text{ yr}^{-1}} \right) \left( \frac{P}{1\%} \right)$$



# What is Required?

## Detecting the Perturbations from Earth-mass Planets

- Sampling rate  $\sim 10$  minutes

$$t_{E,p} = 2\text{hrs} \left( \frac{M_p}{M_E} \right)^{1/2}$$

- Photometric Accuracy  $\sim 1\%$  at  $I \sim 21$

– Signal Magnitude

$$\frac{\Delta F}{F} \approx 1\% \left( \frac{M_p}{M_\oplus} \right) \left( \frac{R_*}{R_\odot} \right)^{-2}$$


– Photometric Uncertainty

$$\sigma = 1\% \left( \frac{D}{2\text{m}} \right)^{-1} \left( \frac{t_{\text{exp}}}{120\text{s}} \right)^{-1/2} 10^{0.2(I-21)}$$

# NextGen $\mu$ Lensing Survey

- Requirements to detect  $\sim 10$  Earth-mass planets per year:
  - Monitor  $\sim 10$  square degrees of the Galactic bulge continuously with  $\sim 10$  minute sampling using 1-2m class telescopes, distributed longitudinally throughout the southern hemisphere.
  - Large FOV (2-4 square degree) cameras needed.

# Expected Results



*A next-generation ground-based  $\mu$ lensing survey can test planet formation by probing planets with  $M > M_{\oplus}$  beyond the snow-line.*

# Spontaneous Generation

- MOA-II (NZ, currently operating)
  - 1.8m telescope, 2.18 sq. degree camera
- OGLE -IV (Chile, 2010)
  - 1.3m telescope, upgrade to 1.4 sq. degree camera
- All that is needed is a 1-2m telescope with a large FOV in South Africa.

*“Recommendation A. II. 1 Increase dramatically the efficiency of a ground-based microlensing network by adding a single 2 meter telescope.”*

# Korean Microlensing Telescope Network (KMTNet)

- A ~30 Billion Won Korean initiative recently been approved by the Korean government (Byeong-Gon Park PI).
  - Three telescopes:
    - South Africa
    - Chile
    - Australia

# Technology

- Ground-based 1-2m, Wide FOV Telescope
  - Several very similar telescopes already operating
    - MOA-II
    - Pan-STARRS-1 - \$20M
- Space-based microlensing mission
  - Requires almost no technology development.
  - Can extensively leverage other missions (Spitzer, NextView, Ikonos, JWST)
  - Can use many components that are demonstrated on orbit or flight qualified.



# Modeling Challenges

- Discovery rate is currently being limited by modeling and manpower.
- Next-generation surveys will exacerbate this problem severely.
  - Order of magnitude larger detection rate.
- Need:
  - More hands (training).
  - Development of automated analysis software.

# Summary

- Ground-based Next-Generation Survey:
  - Support for analysis and software development.
  - Frequency of planets  $>M_{\oplus}$  beyond the snow line.
  - Test planet formation theories.
- **Either:** Space-based Microlensing Mission: +\$300M + launch
  - Complete census of planets with mass greater than Mars and  $a > 0.5$  AU.
  - Sensitivity to all Solar System planet analogs except Mercury.
  - Demographics of planetary systems - tests planet formation theories.
  - Detect “outer” habitable zone (Mars-like orbits) where detection by imaging is easiest.
  - Can find moons and free floating planets.
- **Or:** Joint  $\mu$ lensing/Dark Energy Mission +\$100M—\$200M?
- Total cost to “Exoplanet Community”: **\$100M—\$400M**